



# Photocatalytic hydrogen production over carbon nitride loaded with WS<sub>2</sub> as cocatalyst under visible light



Yidong Hou\*, Yongsheng Zhu, Yan Xu, Xinchun Wang

Research Institute of Photocatalysis, State Key Laboratory Photocatalysis on Energy and Environment, Fuzhou University, Fuzhou 350002, China

## ARTICLE INFO

### Article history:

Received 11 December 2013

Received in revised form 21 February 2014

Accepted 2 March 2014

Available online 12 March 2014

### Keywords:

Carbon nitride

WS<sub>2</sub>

Cocatalyst

Photocatalytic H<sub>2</sub> evolution

## ABSTRACT

Mesoporous graphitic carbon nitride (mpg-CN) was loaded with WS<sub>2</sub> via an impregnation–sulfidation approach. The resultant surface heterojunctions between WS<sub>2</sub> and mpg-CN were evidently confirmed with transmission electron microscopy (TEM) and X-ray photoelectron spectroscopy (XPS). Photocatalytic H<sub>2</sub> production on WS<sub>2</sub>/mpg-CN under visible light ( $\lambda > 420$  nm) shows that WS<sub>2</sub> plays an important role in catalyzing H<sub>2</sub> production. The rate of hydrogen evolution reaches an optimum when the loading amount of WS<sub>2</sub> is about 0.3 at.%. Combined with the physicochemical and electrocatalytic properties of WS<sub>2</sub>/mpg-CN, it can be concluded that the enhanced photocatalytic H<sub>2</sub> production activity on WS<sub>2</sub>/mpg-CN is mainly attributed to the thin layered junctions formed between WS<sub>2</sub> and mpg-CN and the great performance of WS<sub>2</sub> as a cocatalyst in H<sub>2</sub> evolution reaction.

© 2014 Elsevier B.V. All rights reserved.

## 1. Introduction

Since Fujishima and Honda observed a photoelectrochemical cell using Pt–TiO<sub>2</sub> electrodes for hydrogen evolution under the irradiation of ultraviolet (UV) light in 1972 [1], photocatalytic water splitting for hydrogen production, especially based on earth-abundant semiconductors and co-factors, has drawn considerable attentions as a potential way to solve the energy and environment problems [2–4]. So far, many active photocatalysts, including (oxy)nitrides, (oxy)sulfides, and silicides, for H<sub>2</sub> evolution under visible light have been prepared and reported [5–7]. However, most of these photocatalysts under investigation are expensive, unstable, rare, toxic or difficult to prepare. Thus, it is necessary to develop a new type of photocatalyst with a wide visible-light absorption band, high stability in contact with water, non-toxic, and abundant and so on.

Recently, we found that covalent graphitic carbon nitride (g-CN) solid can work as a new type of visible-light and metal-free photocatalyst with suitable electronic structure (conduction band at  $-0.8$  V and valence band at  $1.9$  V vs. RHE) covering redox potentials for water splitting [8,9]. However, like many other photocatalysts, g-CN alone shows a very poor electrocatalytic activity for water splitting, and its photocatalytic performance quite relies on surface cocatalysts [10–14]. The cocatalysts interfaced with the

light-harvesting semiconductor not only facilitate the charge separation and increase the lifetime of the photo-generated electron/hole pair, but also lower the activation barriers for H<sub>2</sub> or O<sub>2</sub> evolution [15,16]. Thus, the use of cocatalysts can result in an increase in overall photocatalytic performance, including activity, selectivity, and stability. Unfortunately, in most cases, the used cocatalysts are mainly the expensive and scarce platinum group metals or their oxides [17]. Therefore, it is one of the most important tasks to replace noble metal by earth abundant elements toward potential large-scale applications [18]. Among many potential candidates, the cheap transition metal based compounds (such as MoS<sub>2</sub>, WS<sub>2</sub>, WC, NiS, Co<sub>3</sub>O<sub>4</sub>) have demonstrated promising activities as the cocatalysts in photocatalytic water splitting reaction [19–23].

In general, the efficiency of a given photocatalytic system is quite dependent on the physical nature of cocatalysts loaded for promoting reductive or/and oxidative catalysis reaction. In particular, the structural characteristics and intrinsic catalytic properties of a cocatalyst are very important [15]. In recent work, we found that MoS<sub>2</sub> is an efficient cocatalyst for the photocatalytic H<sub>2</sub> evolution on mesoporous g-CN (mpg-CN) under visible light as a result of the structural and electronic compatibility between MoS<sub>2</sub> and mpg-CN [13]. Since WS<sub>2</sub> has a similar crystal structure and chemical properties with MoS<sub>2</sub>, we expect that WS<sub>2</sub> could be also used as a promoter similar to MoS<sub>2</sub> for mpg-CN in photocatalytic H<sub>2</sub> production.

Herein, we reported that WS<sub>2</sub>/mpg-CN junctions were obtained through an impregnation–sulfidation process. The detailed structure and property of the resultant WS<sub>2</sub>/mpg-CN were examined.

\* Corresponding author. Tel.: +86 591 83773729; fax: +86 591 83779105.  
E-mail address: [ydhoul@fzu.edu.cn](mailto:ydhoul@fzu.edu.cn) (Y. Hou).

It was found that WS<sub>2</sub> demonstrates similar properties to MoS<sub>2</sub> as a cocatalyst, which can also efficiently promote the photocatalytic activity of mpg-CN in H<sub>2</sub> evolution under visible light irradiation.

## 2. Experiment

### 2.1. Preparation of catalysts

All the chemicals were reagent grade and used without further purification. mpg-CN was synthesized according to the reported procedure [24]. WS<sub>2</sub>/mpg-CN catalysts were prepared via impregnation–sulfidation approach. Certain amounts of tungsten precursor were loaded onto mpg-CN by an impregnation from the solution containing different amounts of (NH<sub>4</sub>)<sub>2</sub>WS<sub>4</sub>. After removing water with a rotary evaporator at 50 °C, the (NH<sub>4</sub>)<sub>2</sub>WS<sub>4</sub>/mpg-CN precursors were heated at 350 °C for 3 h in flowing 10% H<sub>2</sub>S–90% H<sub>2</sub> gas mixture to obtain WS<sub>2</sub>/mpg-CN catalysts. The amount of the WS<sub>2</sub> loaded onto mpg-CN was calculated by assuming that the (NH<sub>4</sub>)<sub>2</sub>WS<sub>4</sub> was completely converted to WS<sub>2</sub>.

### 2.2. Characterization of catalysts

Powder X-ray diffraction (XRD) data were collected using a Bruker D8 Advance X-ray diffractometer (Cu K1 irradiation,  $\lambda = 1.5406 \text{ \AA}$ ). Nitrogen adsorption–desorption isotherms were collected at 77 K using Micromeritics ASAP 2020 equipment. Transmission electron microscopy (TEM) measurements were obtained on a FEI Tencai 20 microscope. X-ray photoelectron spectroscopy (XPS) measurements were performed on a Thermo Scientific ESCA Lab250 spectrometer which consists of a monochromatic Al K $\alpha$  as the X-ray source. All of the binding energies were calibrated by the C1s peak at 284.6 eV. UV–vis diffuse reflectance spectra were collected using a Varian Cary 500 UV–Vis–NIR spectrometer equipped with an integrating sphere, and BaSO<sub>4</sub> was as a reference standard. Photoluminescence (PL) spectra were performed on Edinburgh Analytical Instruments FL/FSTCSPC920 coupled with a time-correlated single-photo-counting system. The electrochemical experiments were conducted on a CHI600E workstation.

### 2.3. Measurement of photocatalytic activity

The photocatalytic reactions were performed in a Pyrex top-irradiation reaction vessel connected to a closed gas circulation and evacuation system. 50 mg of catalyst was suspended in an aqueous solution (100 mL) containing lactic acid (10 vol.%). The reactant solution was degassed several times to remove air prior to irradiation under a Xenon 300 W lamp equipped with various optical cut-off filters to monitor the photocatalytic reaction. The amount of H<sub>2</sub> production was analyzed using an online gas chromatography.

## 3. Results and discussion

Fig. 1 presents the XRD patterns of as-prepared WS<sub>2</sub>/mpg-CN catalysts. It can be seen that mpg-CN has two distinct diffraction peaks at 27.5° and 13.2°, corresponding to the (002) and (100) peaks for graphitic materials, respectively. This is in good agreement with the reported [25]. The positions of the diffraction peaks remain almost unchanged for WS<sub>2</sub>/mpg-CN catalysts obtained via an impregnation–sulfidation approach. Further observation from Fig. 1 reveals that with the increasing of WS<sub>2</sub> content, the diffraction intensity of mpg-CN slightly decreases after loading with WS<sub>2</sub>. However, no characteristic diffraction peaks of WS<sub>2</sub> can be observed. This could be due to the low loading amount and high dispersion of WS<sub>2</sub> in the mpg-CN framework.




Fig. 1. XRD patterns of mpg-CN and WS<sub>2</sub>/mpg-CN.

Nitrogen adsorption–desorption isotherms were used for texture analysis. It can be seen in Fig. 2 that mpg-CN possesses a type H<sub>3</sub> hysteresis loop, implying the presence of mesopores. Textural parameters derived from adsorption–desorption isotherms data are presented in the inserted Table. It can be seen that, comparing with pure mpg-CN, the specific surface area and pore volume of the catalysts decrease largely after loading WS<sub>2</sub>. This could be ascribed to the aggregation of nanoparticles and destruction of porous structure of mpg-CN during the impregnation–sulfidation process.

TEM were used to investigate the morphology of WS<sub>2</sub> and the interfacial structure between WS<sub>2</sub> and mpg-CN. As shown in Fig. 3a, WS<sub>2</sub> slabs with characteristic layered structure instead of particles are found on the surface of WS<sub>2</sub>/mpg-CN. From HRTEM inserted in Fig. 3a, it is easy to find that the (002) plane of WS<sub>2</sub> grows along the surface of mpg-CN to form intimate junctions between WS<sub>2</sub> and mpg-CN. The chemical composition of WS<sub>2</sub>/mpg-CN was then analyzed by using energy-dispersive X-ray spectroscopy (EDX). As displayed in Fig. 3b, the signals assigned to W and S can be easily detected, and which further confirms the existence of WS<sub>2</sub> on the mpg-CN surface. The result of EDX elemental mapping analysis (Fig. S1) verifies that WS<sub>2</sub> is highly dispersed on the surface of mpg-CN. As similar to MoS<sub>2</sub>, WS<sub>2</sub> and graphitic carbon nitride have analogous layered structure and the specific property of Van der Waals interaction may provide advantages in the desirable junction formation. Intimate junctions between WS<sub>2</sub> and mpg-CN are indeed formed by via an impregnation–sulfidation approach.




Fig. 2. Nitrogen adsorption–desorption isotherms of mpg-CN and WS<sub>2</sub>/mpg-CN.




Fig. 3. TEM image (a) and EDX analysis (b) of  $\text{WS}_2/\text{mpg-CN}$ .

Furthermore, the mesoporous structure of mpg-CN facilitates and stabilizes the high dispersion of  $\text{WS}_2$  on the mpg-CN surface and the formation of the thin  $\text{WS}_2/\text{mpg-CN}$  intimate junctions [13]. These junctions ensure effective hydrogen evolution by virtue of the light absorption by mpg-CN combined with a short carrier transport distance, and the exposure of a large external surface area for the catalytic reaction. By taking advantage of electron tunneling effect, the thin-layered  $\text{WS}_2$  highly dispersed on the mpg-CN surface could also offer much higher efficiency than multi-layered  $\text{WS}_2$  for the electron-transporting through the  $\text{WS}_2$  layers to reaction interfaces.

The chemical states of  $\text{WS}_2/\text{mpg-CN}$  were investigated by the XPS technique. As displayed in Fig. 4a, O, N, C, S and W elements are observed in the survey spectrum. Fig. 4b presents the W4f core level spectrum with four peaks at 32.1, 34.1, 34.8 and 36.8 eV. The W4f peaks at 32.1 and 34.1 eV can be ascribed to the  $\text{W}^{4+}$  species, while the other two peaks at higher binding energies can be attributed to  $\text{W}^{6+}$  species [22]. Combined with the result of S2p core level spectrum (Fig. 4c), it confirms that  $\text{WS}_2$  is formed on the surface of mpg-CN. The existence of  $\text{W}^{6+}$  may be explained by that  $(\text{NH}_4)_2\text{WS}_4$  is not stable during the impregnation process, a considerable amount of  $(\text{NH}_4)_2\text{WS}_4$  suffers from oxidation reaction and results in the formation of tungsten oxide. The binding energy of W shifts to a lower value compared to the reported value [26],





Fig. 4. XPS spectra of  $\text{WS}_2/\text{mpg-CN}$ .

indicative of a strong chemical interaction between the W species and the conjugated mpg-CN [23]. Generally, tungsten oxide gradually transforms to  $\text{WS}_2$  with increased sulfidation temperature. Unfortunately, mpg-CN cannot be stabilized at high sulfidation temperature. Thus  $350^\circ\text{C}$  was used as the sulfidation temperature in this study.

Based on the above XPS results analysis, it can be concluded that,  $\text{WS}_2$  and  $\text{WO}_3$  coexist on the surface of mpg-CN. Since the conduction band of  $\text{WO}_3$  situates at a more positive position than the potential of  $\text{H}_2$  evolution [27],  $\text{WO}_3$  is believed to be inactive for photocatalytic  $\text{H}_2$  evolution. Meanwhile, we also annealed the  $(\text{NH}_4)_2\text{WS}_4/\text{mpg-CN}$  in air to prepare  $\text{WO}_3/\text{mpg-CN}$ , and the




**Fig. 5.** UV-vis diffuse reflection spectra (a) and photoluminescence spectra (b) of mpg-CN and WS<sub>2</sub>/mpg-CN.

obtained WO<sub>3</sub>/mpg-CN exhibited negligible activity for photocatalytic hydrogen evolution under visible light. Thus, it is reasonable to consider that the active component for catalytic H<sub>2</sub> evolution is WS<sub>2</sub> rather than WO<sub>3</sub>. Therefore, we assume that WS<sub>2</sub> is the only tungsten species loaded onto mpg-CN in the following sections for simplicity.

Fig. 5a displays the UV-vis diffuse reflection spectra of as-prepared WS<sub>2</sub>/mpg-CN. The abrupt onset of absorption at ~430 nm is assigned to the electronic transition from the valence band to the conduction band in the graphitic carbon nitride semiconductor upon excitation with light [8]. After loading WS<sub>2</sub>, there is a broad absorption shoulder in the spectra. The absorption shoulder corresponds to the optical absorption of WS<sub>2</sub>. Fig. 5b gives the photoluminescence spectra of as-prepared catalysts. Obviously, the photoluminescence of mpg-CN is quenched in the presence of WS<sub>2</sub>. This occurrence is owing to the efficient charge transfer between WS<sub>2</sub> and mpg-CN, thus leading to an improvement in the separation efficiency of the photo-generated carriers.

No H<sub>2</sub> was detected when WS<sub>2</sub> was supported on inert mesoporous SiO<sub>2</sub> (SBA-15), thus implying that WS<sub>2</sub> alone is not active for photocatalytic H<sub>2</sub> evolution in a wireless system. Pure mpg-CN also showed negligible activity. Fig. 6a shows the rate of H<sub>2</sub> evolution on the WS<sub>2</sub>/mpg-CN catalysts. There is a plot of the variation in their activities, showing that the rate of hydrogen evolution increases initially and then reaches an optimum when the amount of the loading of WS<sub>2</sub> is about 0.3 at.%. Further loading of WS<sub>2</sub> on mpg-CN leads to a decrease in the H<sub>2</sub> evolution rate. The decrease in the



**Fig. 6.** (a) Rate of H<sub>2</sub> evolution on mpg-CN with loading different amounts of WS<sub>2</sub> under visible light ( $\lambda > 420$  nm), (b) cyclic runs for the photocatalytic H<sub>2</sub> production on WS<sub>2</sub>/mpg-CN.

activity of the catalysts with a heavy loading of WS<sub>2</sub> could be due to the shading effect, which can seriously block the absorption of the incident light by mpg-CN. This hypothesis is strongly supported by the UV-vis diffuse reflection spectra.


We also investigated the rate of photocatalytic hydrogen evolution on WS<sub>2</sub>/mpg-CN as a function of cut-off wavelength of the incident light. Fig. S2 reveals that the wavelength dependence of hydrogen evolution rate of WS<sub>2</sub>/mpg-CN corresponds well to its optical absorption. This observation supports that the hydrogen evolution process is primarily induced by light-excitation of carbon nitride polymer.

In addition to the photocatalytic activity, the stability of a photocatalyst is also important in practical application. In order to study the stability of WS<sub>2</sub>/mpg-CN, the cycling hydrogen evolution experiments were performed. As shown in Fig. 6b, the rate of H<sub>2</sub> evolution on WS<sub>2</sub>/mpg-CN decreased with every reaction cycle. To reveal the reason for the deactivation during activity test, we did the comparisons of XRD and XPS analysis on the fresh and used WS<sub>2</sub>/mpg-CN. The XRD analysis result shows that the crystal structure of WS<sub>2</sub>/mpg-CN does not change after the photocatalytic test (Fig. S3). The fresh and used photocatalyst give similar N1s and C1s core level spectra shown in Fig. S4, this further confirms the high stability of mpg-CN. However, significant changes are observed in W4f and S2s core level spectra (Fig. S4). The data (Table 1) obtained after deconvolution of these peaks reveal that the atomic ratio of W<sup>6+</sup>/N increases and that of S<sup>2-</sup>/N decreases for the catalyst after test. This means that the oxidation/corrosion of WS<sub>2</sub> by the



**Table 1**  
Values of W and S species to N in the fresh and used WS<sub>2</sub>/mpg-CN.

Sample	W/N (at.%)			S/N(at.%)		
	W <sup>4+</sup>	W <sup>6+</sup>	W (sum)	S <sup>2-</sup>	S <sup>6+</sup>	S (sum)
WS <sub>2</sub> /mpg-CN (fresh)	0.13	0.21	0.34	0.68	0.25	0.93
WS <sub>2</sub> /mpg-CN (used)	0	0.25	0.25	0	0	0



**Fig. 7.** Polarization curves of mpg-CN/FTO and WS<sub>2</sub>/mpg-CN/FTO electrodes.

photo-generated holes occurs on WS<sub>2</sub>/mpg-CN catalyst, accounting for the deactivation during the cycling runs. Of course, more work is required to improve the activity and particularly stability of WS<sub>2</sub>/mpg-CN for potential application, for example to release charge built-up by using oxidative cofactors to promote the transfer of the holes from the bulk to the surface.


It is well regarded that the reduction of protons to H<sub>2</sub> is a very important step in the photocatalytic H<sub>2</sub> production reaction. It has been reported that, the catalytic characteristics of MoS<sub>2</sub> for H<sub>2</sub> evolution is a very important factor for the enhanced activity in photocatalytic water splitting system [13,28]. In this study, electrochemical measurements were conducted to investigate the role of WS<sub>2</sub>. Fig. 7 shows the polarization curve of the WS<sub>2</sub>/mpg-CN/FTO electrode together with that of mpg-CN/FTO as a control. Cathodic current observed in the potential range from −0.9 to −1.4 V vs.

Ag/AgCl corresponds to the reduction of water to H<sub>2</sub>. In comparison with mpg-CN/FTO electrode, WS<sub>2</sub>/mpg-CN/FTO electrode shows much enhanced cathodic current. Apparently, WS<sub>2</sub> is a good catalyst that can efficiently catalyze the H<sub>2</sub> evolution, thus enhances the rate of H<sub>2</sub> evolution on WS<sub>2</sub>/mpg-CN. Moreover, because of the anisotropic conductivity of WS<sub>2</sub>, like MoS<sub>2</sub> [29], thin-layered WS<sub>2</sub> can lessen the poor charge transport from layer to layer and shorten the electron transport time and distance, thus improves the efficiency in the utilization of photo-generated electrons for hydrogen production.

On the basis of the above results, the mechanism for the photocatalytic H<sub>2</sub> production on WS<sub>2</sub>/mpg-CN is proposed. As shown in Scheme 1, upon photo-excitation, electrons and holes are generated in the conduction band (CB) and valence band (VB) of mpg-CN, respectively. The photo-generated carriers move toward the catalyst surface, the holes oxidize the lactic acid and the electrons reduce protons to H<sub>2</sub>. However, because mpg-CN itself is not active for H<sub>2</sub> evolution, the photo-generated electrons and holes are likely to recombine, resulting in the low photocatalytic activity. The presence of WS<sub>2</sub> facilitates the transfer of the photo-generated electrons from mpg-CN to WS<sub>2</sub>, and protons can be efficiently reduced to produce H<sub>2</sub> because WS<sub>2</sub> can serve as an active site for hydrogen evolution. Moreover, the formed thin-layered junctions promote the electron transfer from mpg-CN to WS<sub>2</sub>. These cofactors are supposed to be the main reasons why the WS<sub>2</sub>/mpg-CN exhibited enhanced activity of hydrogen evolution.

#### 4. Conclusion

The H<sub>2</sub> production performance of mpg-CN under visible light is significantly improved by loading WS<sub>2</sub>. The rate of hydrogen evolution reaches an optimum when the amount of WS<sub>2</sub> loaded is about 0.3 at.%. Similar to MoS<sub>2</sub>, the geometric similarity in the layered



**Scheme 1.** Proposed reaction mechanism for photocatalytic H<sub>2</sub> evolution on WS<sub>2</sub>/mpg-CN.

structure of WS<sub>2</sub> and mpg-CN and the mesoporous structure of mpg-CN facilitate the planar growth of WS<sub>2</sub> on the mpg-CN surface and the formation of the thin planar WS<sub>2</sub>/mpg-CN interface. These characteristics are supposed to be mainly responsible for the enhanced photocatalytic activity of WS<sub>2</sub>/mpg-CN catalysts. This work demonstrates a possibility for the use of WS<sub>2</sub> as a substitute for noble metals as cocatalyst in the photocatalytic H<sub>2</sub> production. It also provides useful information in design and fabricating effective interfacial junctions between cocatalysts and semiconductors.

### Acknowledgements

The work was supported by 973 Program (2013CB632405), NSFC (21203030 and 21033003), Specialized Research Fund for the Doctoral Program of Higher Education Program (20113514120001) and Program for New Century Excellent Talents in University of Fujian Province (JA13020).

### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.apcatb.2014.03.002>.

### References

- [1] A. Fujishima, K. Honda, *Nature* 238 (1972) 37–38.
- [2] J.A. Turner, *Science* 305 (2004) 972–974.
- [3] N.S. Lewis, D.G. Nocera, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 15729–15735.
- [4] M. Pelaez, N.T. Nolan, S.C. Pillai, M.K. Seery, P. Falaras, et al., *Appl. Catal. B Environ.* 125 (2012) 331–349.
- [5] A. Kudo, *Int. J. Hydrogen Energy* 31 (2006) 197–202.
- [6] F.E. Osterloh, *Chem. Mater.* 20 (2008) 35–54.
- [7] X.B. Chen, S.H. Shen, L.J. Guo, S.S. Mao, *Chem. Rev.* 110 (2010) 6503–6570.
- [8] X.C. Wang, K. Maeda, A. Thomas, K. Takanabe, G. Xin, et al., *Nat. Mater.* 8 (2009) 76–80.
- [9] J.S. Zhang, X.F. Chen, K. Takanabe, K. Maeda, K. Domen, et al., *Angew. Chem. Int. Ed.* 49 (2010) 441–444.
- [10] K. Maeda, X.C. Wang, Y. Nishihara, D.L. Lu, M. Antonietti, et al., *J. Phys. Chem. C* 113 (2009) 4940–4947.
- [11] Y. Di, X.C. Wang, A. Thomas, M. Antonietti, *ChemCatChem* 2 (2010) 834–838.
- [12] A.T. Garcia-Esparza, D. Cha, Y. Ou, J. Kubota, K. Domen, et al., *ChemSusChem* 6 (2013) 168–181.
- [13] Y.D. Hou, A.B. Laursen, J.S. Zhang, G.G. Zhang, Y.S. Zhu, et al., *Angew. Chem. Int. Ed.* 52 (2013) 3621–3625.
- [14] L. Ge, C.C. Han, X.L. Xiao, L.L. Guo, *Appl. Catal. B Environ.* 142–143 (2013) 414–422.
- [15] K. Maeda, K. Domen, *J. Phys. Chem. Lett.* 1 (2010) 2655–2661.
- [16] J.H. Yang, D.G. Wang, H.X. Han, C. Li, *Acc. Chem. Res.* 46 (2013) 1900–1909.
- [17] M.G. Walter, E.L. Warren, J.R. Mc-Kone, S.W. Boettcher, Q.X. Mi, et al., *Chem. Rev.* 110 (2010) 6446–6473.
- [18] P.C.K. Vesborg, T.F. Jaramillo, *RSC Adv.* 2 (2012) 7933–7947.
- [19] X. Zong, H.J. Yan, G.P. Wu, G.J. Ma, F.Y. Wen, et al., *J. Am. Chem. Soc.* 130 (2008) 7176–7177.
- [20] J.S. Jang, D.J. Ham, N. Lakshminarasimhan, W. Choi, J.S. Lee, *Appl. Catal. A Gen.* 346 (2008) 149–154.
- [21] W. Zhang, Y. Wang, Z. Wang, Z. Zhong, R. Xu, *Chem. Commun.* (2010) 7631–7633.
- [22] X. Zong, J.F. Han, G.J. Ma, H.J. Yan, G.P. Wu, et al., *J. Phys. Chem. C* 115 (2011) 12202–12208.
- [23] J.S. Zhang, M. Grzelczak, Y.D. Hou, K. Maeda, K. Domen, et al., *Chem. Sci.* 3 (2012) 443–446.
- [24] X.C. Wang, K. Maeda, X.F. Chen, K. Takanabe, K. Domen, et al., *J. Am. Chem. Soc.* 131 (2009) 1680–1681.
- [25] A. Thomas, A. Fischer, F. Goettmann, M. Antonietti, J. Muller, et al., *J. Mater. Chem.* 18 (2008) 4893–4908.
- [26] A. Di-Paola, L. Palmisano, A.M. Venezia, V. Augugliaro, *J. Phys. Chem. B* 103 (1999) 8236–8244.
- [27] Y. Xu, M.A.A. Schoonen, *Am. Miner.* 85 (2000) 514–523.
- [28] X. Zong, G.P. Wu, H.J. Yan, G.J. Ma, J.Y. Shi, et al., *J. Phys. Chem. C* 114 (2010) 1963–1968.
- [29] H. Tributsch, J.C. Bennett, *J. Electroanal. Chem.* 81 (1977) 97–111.